

Limits on the ν_τ mass from nucleosynthesis in the presence of annihilation into majorons¹

J. C. Romão²

*Instituto Superior Técnico, Departamento de Física
A. Rovisco Pais, 1 1096 Lisboa Codex, Portugal*

Contribution to the International Workshop
New Worlds in Astroparticle Physics
Faro, Portugal, 8-10 September 1996.

Abstract

We show that in the presence of sufficiently strong ν_τ annihilations into majorons, the primordial nucleosynthesis constraints can not rule out ν_τ masses in the MeV range.

¹Work done in collaboration with A.D. Dolgov (Denmark), S. Pastor and J.W.F Valle (IFIC, València)

²E-mail fromao@alfa.ist.utl.pt

1 Introduction

Despite great experimental efforts the tau-neutrino still remains as the only one that can have a mass in the MeV range. The present experimental limit on its mass is [1]

$$m_{\nu_\tau} < 23 \text{ MeV} \quad (1)$$

Further progress will have to wait for the tau-charm or B-factories. On the other hand, many particle physics models of massive neutrinos lead to a tau neutrino with mass in the MeV range. Moreover such a neutrino may have interesting cosmological implications. It is therefore important to examine in more detail the cosmological constraints.

The first comes from the critical density argument. From the contribution of stable ν_τ to the present relic density one gets $m_{\nu_\tau} < 92\Omega h^2 \text{ eV}$, where h is related to the Hubble constant through $h = H_0/(100 \text{ kms}^{-1} \text{ Mpc}^{-1})$. This means that a massive ν_τ with mass in the MeV range must be unstable with lifetimes smaller than the age of the Universe. However, it has been shown that in many particle physics models where neutrinos acquire their mass by the spontaneous breaking of a global lepton number symmetry[2] this limit can be avoided due to the existence of fast ν_τ decays.

The second constraint comes from primordial nucleosynthesis considerations. In the standard model, these rule out ν_τ masses in the range[3, 4]

$$0.5 \text{ MeV} < m_{\nu_\tau} < 35 \text{ MeV} \quad (2)$$

It is possible to weaken the constraints on ν_τ of Eq. (2), by adding new interactions beyond the standard ones. One may either consider that the ν_τ 's are unstable during nucleosynthesis[5] or that they possess new channels of annihilation beyond the standard ones. It is this last possibility that we investigate in the case where the ν_τ 's can annihilate to majorons (J).

2 Evolution of ν_τ number density

2.1 Before weak decoupling

We will assume that the massive Majorana ν_τ 's are stable during the nucleosynthesis epoch. They interact with leptons via the standard weak interactions $\nu_\tau \bar{\nu}_\tau \leftrightarrow \nu_0 \bar{\nu}_0, e^+ e^-$. Moreover the ν_τ 's annihilate to majorons $\nu_\tau \nu_\tau \rightarrow JJ$ via the diagrams shown in Fig. (1). The t-channel diagram is present in all Majoron models, while the strength of the s-channel scalar exchange diagram is somewhat model-dependent. For the range of g values relevant for our purposes ($g \gtrsim 10^{-5}$) it is quite reasonable to neglect the s-channel process, especially in Majoron models where the breakdown of the global lepton number symmetry happens at the electroweak scale or higher, like the simplest seesaw model [6]. We have shown[7] that for models with low




Figure 1: Feynman diagrams for annihilations of tau neutrinos into Majorons.

scale breaking of lepton number, such as given in ref. [8, 9] the bounds obtained by neglecting the s-channel process could be relaxed by at most a factor ~ 2 or so. Therefore in what follows we will consider that the ν_τ 's annihilate to majorons $\nu_\tau \nu_\tau \rightarrow JJ$ via the diagonal coupling

$$\mathcal{L}_{\nu_\tau \nu_\tau J} = i \frac{1}{2} g J \overline{\nu_\tau} \gamma_5 \nu_\tau \quad (3)$$

The evolution of the number densities of ν_τ (n_τ) and majorons (n_J) is given by the solution of a set of Boltzmann differential equations

$$\dot{n}_\tau + 3Hn_\tau = - \sum_{i=e, \nu_0} \langle \sigma_i v \rangle (n_\tau^2 - (n_\tau^{eq})^2) - \langle \sigma_J v \rangle (n_\tau^2 - (n_\tau^{eq})^2 \frac{n_J^2}{(n_J^{eq})^2}) \equiv S_{\nu_\tau} \quad (4)$$

$$\dot{n}_J + 3Hn_J = \langle \sigma_J v \rangle (n_\tau^2 - (n_\tau^{eq})^2 \frac{n_J^2}{(n_J^{eq})^2}) \equiv S_J \quad (5)$$

where $\langle \sigma_i v \rangle$ are the thermally averaged cross sections times the ν_τ relative velocity v . The cross sections are for annihilation to majorons

$$\sigma_J(\eta) = \frac{g^4}{128\pi} \frac{1-\eta}{m_{\nu_\tau}^2 \eta} \left[\ln \left(\frac{1+\sqrt{\eta}}{1-\sqrt{\eta}} \right) - 2\sqrt{\eta} \right] \quad (6)$$

and for annihilation into *massless* fermions

$$\sigma_i(\eta) = \frac{32G_F^2}{3\pi} \frac{m_{\nu_\tau}^2 \sqrt{\eta}}{1-\eta} (b_{Li}^2 + b_{Ri}^2) \quad (7)$$

where $b_L^2 + b_R^2 = 1/2$ for $i = \nu_0$ and $b_L^2 + b_R^2 = 2 \left[(-1/2 + \sin^2 \theta_W)^2 + (\sin^2 \theta_W)^2 \right] \simeq 0.25$ for $i = e$. In Eq. (5) we assumed Boltzmann statistics and $n_i = n_i^{eq}$ for $i = e, \nu_0$.

Now let us describe our calculations. First we normalized the number densities to the number density of massless neutrinos, $n_0 \simeq 0.181T^3$, introducing the quantities

$r_\alpha \equiv n_\alpha/n_0$, where $\alpha = \nu_\tau, J$, and the corresponding equilibrium functions r_α^{eq} . Then from Eq. (5) we get

$$\frac{dr_\alpha}{dT} = \left(\frac{S_\alpha}{n_0} - 3Hr_\alpha \right) \frac{1}{\dot{T}} - \frac{3}{T} r_\alpha \quad (8)$$

On the other hand, the time derivative of the temperature is obtained from Einstein's equation

$$\dot{\rho} = -3H(\rho + P) \quad (9)$$

where ρ is the total energy density and P is the pressure.

2.2 Past weak decoupling

Once the ν_τ 's decouple from the standard weak interactions, they remain in contact only with the majorons. Then one has two different plasmas, one formed by ν_τ 's and J 's and the other by the rest of particles, each with its own temperature, denoted by T and T_γ , respectively. We assume that the photon temperature evolves in the usual way, $\dot{y} = Hy$. The evolution equations of the ν_τ and J number densities are now simplified versions of Eq. (5), because we $S_{\nu_\tau} = -S_J$,

$$\begin{aligned} \dot{n}_\tau + 3Hn_\tau &= -S_J \\ \dot{n}_J + 3Hn_J &= S_J \end{aligned} \quad (10)$$

To solve these equations one must find a relation between T and T_γ . This is obtained using Einstein's equation for the $\nu_\tau + J$ plasma.

In order to determine the final ν_τ frozen density which will be relevant during nucleosynthesis we have to solve numerically the appropriate set of coupled differential equations for each pair of values (m_{ν_τ}, g) . Before weak decoupling these are Eqs. (5) e (9) while after decoupling one should use Eq. (10). The initial conditions are, for sufficiently high temperatures, $r_{\nu_\tau} = r_{\nu_\tau}^{eq}$, $r_J = r_J^{eq}$ and $T_{\nu_\tau} = T_J = T_{\nu_0}$.

3 Nucleosynthesis constraints

The value of $r_{\nu_\tau}(m_{\nu_\tau}, g)$ is used to estimate the variation of the total energy density $\rho_{tot} = \rho_R + \rho_{\nu_\tau}$. In ρ_R all relativistic species are taken in account, including the majorons and two massless neutrinos, whereas ρ_{ν_τ} is the energy density of the massive ν_τ 's.

In order to compare with the standard model situation, we can now express the effect of the ν_τ mass and of the annihilation to majorons in terms of an effective number of massless neutrino species, N_{eq} , which we calculate for each value of $r_{\nu_\tau}(m_{\nu_\tau}, g)$. To do this, we first numerically calculate the evolution of the neutron fraction, r_n , by varying the value of N_{eq} [10]. Then we incorporate ρ_{tot} in this numerical code and perform the integration for each pair of (m_{ν_τ}, g) values. Comparing




Figure 2: Effective number of massless neutrinos equivalent to the contribution of heavy ν_τ 's with different values of g in units of 10^{-5} . For comparison, the dashed line corresponds to the standard model case when $g = 0$.

the r_n obtained in each case at $T_\gamma \simeq 0.065 \text{ MeV}$ (the moment when practically all neutrons are wound up in ${}^4\text{He}$), we can relate (m_{ν_τ}, g) and N_{eq} .

The results are shown in Fig. (2). One can see that for fixed N_{eq} , a wide range of tau neutrino masses is allowed for large enough coupling constants. In fact all masses below 23 MeV are allowed provided the coupling constant g exceeds a value a few times 10^{-4} . For comparison, the dashed line corresponds to the case when $g = 0$ and no majorons are present. These results can also be expressed in the $m_{\nu_\tau} - g$ plane, as shown in Fig. (3).

4 Majoron models

There has been a variety of majoron models proposed in the literature[11]. They are attractive extensions of the standard model where neutrinos acquire mass by virtue of the spontaneous violation of a global lepton number symmetry. Apart from a phenomenological interest of their own, majoron models offer the possibility of loosening the cosmological limits on the neutrino masses, either because neutrinos decay or annihilate to majorons.

We have seen above that the restrictions imposed by primordial nucleosynthesis upon a heavy tau neutrino disappear for sufficiently large values of the $\nu_\tau \nu_\tau$ majoron coupling g . Different models imply different expectations for the coupling g and for the relation between g and the ν_τ mass. Just to give a concrete example let us consider the supersymmetric models with spontaneous violation of R parity[9]. In these models we get the relation between g and m_{ν_τ} depicted in Fig. (4). The




Figure 3: The values of $g(m_{\nu_\tau})$ above each line would be allowed by nucleosynthesis if one adopts the $N_{eq}^{max} = 3, 3.4, 3.8, 4.2$ (from top to bottom).

different curves correspond to different values of the parameter v_R of the model. We see that we can obtain for g values of the order of a few times 10^{-4} , as required in our nucleosynthesis analysis.

5 Conclusions

We have determined the restrictions imposed by primordial nucleosynthesis upon a heavy tau neutrino in the presence of sufficiently strong ν_τ annihilations into majorons. We have shown that for values of the ν_τ ν_τ majoron coupling in excess of 10^{-4} all the masses up to the experimental limit of 23 MeV can be allowed. We have shown that in a supersymmetric model with spontaneous violation of R parity such values can be obtained.

References

- [1] D. Buskulic et al., *Phys. Lett.* **B349** (1995) 585.
- [2] For a recent review see J.W.F. Valle, in *Physics Beyond the Standard Model*, lectures given at the *VII Jorge André Swieca Summer School* (Rio de Janeiro, February 1995) and at *V Taller Latinoamericano de Fenomenologia de las Interacciones Fundamentales* (Puebla, Mexico, October 1995); [hep-ph/9603307](#).
- [3] E. W. Kolb, M.S. Turner, A. Chakravorty and D. N. Schramm, *Phys. Rev. Lett.* **67** (1991) 533.




Figure 4: Expected values of m_{ν_τ} and g in model of ref [9]

- [4] A. D. Dolgov and I. Z. Rothstein, *Phys. Rev. Lett.* **71** (1993) 476
- [5] M. Kawasaki, G. Steigman and H.-S. Kang, *Nucl. Phys.* **B402** (1993) 323, *Nucl. Phys.* **B419** (1994) 105; S. Dodelson, G. Gyuk and M.S. Turner, *Phys. Rev.* **D49** (1994) 5068.
- [6] Y. Chikashige, R. Mohapatra and R. Peccei, *Phys. Rev. Lett.* **45** (1980) 1926.
- [7] A. D. Dolgov, S. Pastor, J.C. Romão and J. W. F. Valle, [hep-ph/9612311](#).
- [8] M.C. González-García and J.W.F. Valle, *Phys. Lett.* **B216** (1989) 360; M.C. González-García, A. Santamaria and J.W.F. Valle, *Nucl. Phys.* **B342** (1990) 108; A. Joshipura and J.W.F. Valle, *Nucl. Phys.* **B397** (1993) 105; J.T. Peltoniemi, and J.W.F. Valle, *Phys. Lett.* **B304** (1993) 147; Z. Berezhiani, A.Yu. Smirnov and J.W.F. Valle, *Phys. Lett.* **B291** (1992) 99; J.C. Romão, F. de Campos, and J.W.F. Valle, *Phys. Lett.* **B292** (1992) 329.
- [9] A. Masiero and J. W. F. Valle, *Phys. Lett.* **B251** (1990) 273; J. C. Romão, C. A. Santos and J. W. F. Valle, *Phys. Lett.* **B288** (1992) 311.
- [10] D.A. Dicus *et al.*, , *Phys. Rev.* **D26** (1982) 2694.
- [11] For recent reviews see J. W. F. Valle, *Gauge Theories and the Physics of Neutrino Mass*, *Prog. Part. Nucl. Phys.* **26** (1991) 91-171 (ed. A. Faessler)